

# EGO-1, a Putative RNA-Dependent RNA Polymerase, Is Required for Heterochromatin Assembly on Unpaired DNA during *C. elegans* Meiosis

Eleanor M. Maine,<sup>1,\*</sup> Jessica Hauth,<sup>2</sup> Thomas Ratliff,<sup>2</sup> Valarie E. Vought,<sup>1</sup> Xingyu She,<sup>1</sup> and William G. Kelly<sup>2,\*</sup>

<sup>1</sup>Department of Biology  
Syracuse University  
Syracuse, New York 13244

<sup>2</sup>Biology Department  
Emory University  
Atlanta, Georgia 30322

## Summary

During meiosis in *C. elegans*, unpaired chromosomes and chromosomal regions accumulate high levels of histone H3 lysine 9 dimethylation (H3K9me2), a modification associated with facultative heterochromatin assembly and the resulting transcriptional silencing [1, 2]. Meiotic silencing of unpaired DNA may be a widely conserved genome defense mechanism [3–5]. The mechanisms of meiotic silencing remain unclear, although both transcriptional and posttranscriptional processes are implicated [3–5]. Cellular RNA-dependent RNA polymerases (RdRPs) function in development and RNA-mediated silencing in many species [3, 6, 7] and in heterochromatin assembly in *S. pombe* [3, 8]. There are four *C. elegans* RdRPs, including two with known germline functions. EGO-1 is required for fertility and robust germline RNAi [9–11]. RRF-3 acts genetically to repress RNAi and is required for normal meiosis and spermatogenesis at elevated temperatures [12] (S. L'Hernault, personal communication). Among *C. elegans* RdRPs, we find that only EGO-1 is required for H3K9me2 enrichment on unpaired chromosomal regions during meiosis. This H3K9me2 enrichment does not require Dicer or Drosha nuclease or any of several other proteins required for RNAi. *ego-1* interacts genetically with *him-17*, another regulator of chromatin and meiosis [13], to promote germline development. We conclude that EGO-1 is an essential component of meiotic silencing in *C. elegans*.

## Results and Discussion

### EGO-1 and RRF-3 Activity Regulate Meiotic H3K9me2 Accumulation

Since RdRP activity is required for meiosis and meiotic silencing in *N. crassa* and for heterochromatin assembly in *S. pombe*, we investigated the effect of RdRP mutations on meiotic heterochromatin assembly in *C. elegans* using anti-H3K9me2 antibodies. (Details of strains used throughout this report are described in the [Supplemental Experimental Procedures](#).) We first investigated *ego-1* mutants. We evaluated three types of unpaired chromosomes that become enriched for H3K9me2 during meiosis [1, 2]: (1) the single male X chromosome,

(2) poorly paired X chromosomes that occur in *him-8* (high incidence of males) mutants, and (3) an unpaired autosomal fragment, *sDp3*. Each type of unpaired DNA became enriched in H3K9me2 during meiosis in wild-type (*ego-1(+)*) animals and *ego-1(0)/+* heterozygotes, but failed to do so in *ego-1(0)* null mutants (Figures 1A–1C). H3K9me2 accumulation was also defective in males carrying an *ego-1* point mutation in a conserved residue in the RdRP catalytic domain in *trans* to a null allele (designated *ego-1(RdRP/null)*) (data not shown).


In wild-type hermaphrodites, H3K9me2 staining is briefly elevated on autosomes in mid-to-late pachytene and diplotene stages. This transient accumulation is excluded from paired X chromosomes, which undergo activation during oogenesis [1, 2]. In *ego-1(0)* hermaphrodites, we still observed a low level of H3K9me2 accumulation on autosomes (Figure 1B; data not shown). We conclude that EGO-1 activity is required for H3K9me2 enrichment on unpaired chromosomal regions during pachytene stage of meiosis, but not for accumulation on autosomes in later stages.

*rrf-3(0)* mutants have a temperature-sensitive germline phenotype [12]; therefore, we assayed H3K9me2 targeting in *rrf-3(0)* mutants raised at restrictive (25°C) temperature. We consistently observed H3K9me2 targeting to the male X in these animals (see Figure S1 in the [Supplemental Data](#) available with this article online). Strikingly, H3K9me2 staining persisted in *rrf-3(0)* mutants substantially longer than in wild-type, remaining strong in primary spermatocytes (Figure S1). H3K9me2 is normally depleted from chromatin by the time primary spermatocyte nuclei begin to visibly condense and before other modifications are extensively depleted from the autosomes (Figure 1A; Figure S1; [2]). The temperature-sensitive fertility defects in the *rrf-3(0)* mutants may correlate with the aberrant chromatin regulation that we observe in these animals. H3K9me2 accumulation during meiosis was normal in *rrf-1(0)* and *rrf-2(0)* males (data not shown).

### Regulation of Repetitive DNA Arrays in the *ego-1* Mutant Germline

*C. elegans* transgenes are often generated as multicopy, extrachromosomal arrays [14]. Expression of such transgenes is typically detected in the soma, but not in the germline [2, 15]. Germline-silenced arrays accumulate a high density of H3K9me2 during pachytene stage (similar to the male X) and do not accumulate “active” chromatin modifications, e.g., H3 methylated on lysine 4 (H3K4me) [2]. We investigated whether EGO-1 is required for H3K9me2 accumulation on a repetitive extrachromosomal array, *ccEx7291*, and a repetitive array fused to a chromosomal fragment, *svDp1*. Strong H3K9me2 staining of both arrays was seen in wild-type and *ego-1(0)/+* animals, but not in *ego-1(0)* animals (Figure 2A). Therefore, EGO-1 activity is critical for targeting H3K9me2 to unsynapsed, repetitive arrays. We note that these transgenes are not likely to be fully paired with any partner during meiosis.

\*Correspondence: emmaine@syr.edu (E.M.M.); william.kelly@emory.edu (W.G.K.)



**Figure 1. EGO-1 Activity Is Required for H3K9me2 Accumulation on Unpaired Chromosomes and Chromosomal Fragments during Meiosis**  
Each panel shows meiotic nuclei stained with DAPI to visualize DNA or polyclonal anti-H3K9me2 antibody. *ego-1* null alleles are used in all cases (see Supplemental Experimental Procedures). Photographs are oriented with distal tissue to the left.

(A) Wild-type (wt) male pachytene cells contain a bright focus of H3K9me2 staining associated with the condensed X chromosome. Weaker staining is visible on autosomes. H3K9me2 staining rapidly decreases as cells exit pachytene and start to condense as primary spermatocytes (also see Figure S1). In *ego-1* mutant males, the condensed X chromosome is distinct in DAPI-stained preparations but fails to accumulate a bright focus of H3K9me2 staining (arrows). This result is consistent with previous work showing that condensation of the male X can occur in the absence of H3K9me2 enrichment [1]. Weak H3K9me2 staining is visible on all chromosomes. We note that occasional *ego-1* nuclei had one or more brighter foci of staining that did not seem to localize to the X, suggesting that EGO-1 activity may also prevent inappropriate H3K9me2 accumulation on autosomes.

(B) *him-8* and *ego-1/+;him-8* pachytene cells have two bright foci of H3K9me2 staining, corresponding to the unpaired regions of the X chromosomes (arrows). In some nuclei, these regions are not adjacent. Paired autosomes have weaker staining, as in wild-type. In contrast, the bright foci of H3K9me2 staining are not evident in *ego-1;him-8* pachytene cells.


(C) *ego-1/+;sDp3* animals have 1–2 bright foci of H3K9me2 staining in pachytene cells (arrows). Two foci result when partial pairing occurs between the left end of *sDp3* and the left end of an intact chromosome III, leaving a portion of *sDp3* and a portion of the other chromosome III unpaired. In contrast, *ego-1;sDp3* pachytene cells lack the bright H3K9me2 foci on both *sDp3* and the unpaired region of the intact chromosome III.

(D) Both *dcr-1/+* and *dcr-1* males have a bright focus of H3K9me2 staining in pachytene cells that corresponds to the tightly condensed X chromosome (arrows). Weaker staining is associated with autosomes.

*ccEx7291* carries a GFP reporter, *let-858::gfp*. *let-858::gfp* has been used to detect genes that participate in germline silencing, including components of both transcriptional and posttranscriptional mechanisms (TGS and PTGS, respectively) [15–18]. Although *ccEx7291* did not accumulate H3K9me2 in the *ego-1(0)* mutant, we did not detect germline expression of LET-858::GFP (data not shown). We tested other GFP-tagged transgenes and also found that they remained silenced in *ego-1(0)* germ cells (data not shown).

Our findings may reflect the regulation of multicopy arrays by multiple mechanisms. We examined H3K9me2 accumulation on *ccEx7291* in *rde-2* (RNAi defective)

mutants, which are defective in germline PTGS and transgene silencing [19]. Although LET-858::GFP was expressed (at a low level), *ccEx7291* accumulated normal H3K9me2 levels (Figure S2). H3K9me2 accumulation is likely just one of several mechanisms that limit expression of repetitive DNA in the germline. Weak germline expression of extrachromosomal arrays is seen in certain PTGS mutants, including *dcr-1* and *rde-2*, and also in *mes* mutants, which are defective in regulation of the silencing mark, H3K27me [15, 19–23; this report]. We may fail to see accumulation of LET-858::GFP in *ego-1* mutant germlines because these other repressive mechanisms remain active.



**Figure 2. H3K9me2 Accumulation on Extrachromosomal, Multicopy Transgenes Is Defective in *ego-1* but Not in *dcr-1* Mutants**  
Each panel shows a portion of the meiotic germline from a strain carrying the repetitive extrachromosomal array, *ex7291*. (A) *ex7291* is visible in DAPI-stained preparations of both *ego-1/+* and *ego-1* tissue (see insets; arrows indicate extrachromosomal array). A bright focus of H3K9me2 staining, corresponding to *ex7291*, is visible in *ego-1/+* (arrows) but not *ego-1* tissue. (B) A bright focus of H3K9me2 staining, corresponding to *ex7291*, is visible in both *dcr-1/+* and *dcr-1* tissue. Scale bar equals 20  $\mu$ m; inset scale bar equals 5  $\mu$ m.

### H3K9me2 Accumulation in *dcr-1/Dicer* and *drsh-1/Drosha* Mutants

Dicer endonuclease functions in heterochromatin assembly in many species and may act in the biogenesis of short RNAs that recruit the chromatin-modifying machinery to target loci [3, 8, 24, 25]. We investigated whether *C. elegans* DCR-1/Dicer activity is required for H3K9me2 accumulation on unpaired DNA during meiosis. H3K9me2 accumulation appeared normal on the X chromosome in *dcr-1(0)* adult males, on partially paired X chromosomes in *dcr-1(0);him-8* adult hermaphrodites, and on autosomes in late pachytene-diplotene stage in *dcr-1(0)* hermaphrodites (Figure 1D; data not shown). Moreover, H3K9me2 accumulated normally on *ccEx7291* in a *dcr-1(0)* background (Figure 2B). Therefore, DCR-1 does not seem to be required for H3K9me2 accumulation during meiosis. One caveat is that these *dcr-1(0)* mutants were derived from *dcr-1(0)/+* heterozygous mothers and therefore had inherited maternal product. Embryos lacking both maternal and zygotic DCR-1 arrest during early development and cannot be assayed for defects in germline chromatin (data not shown).

We evaluated the importance of another RNase III-type nuclease, Drosha [26]. H3K9me2 staining was normal in *drsh-1(0)* males and *drsh-1(0);him-8* hermaphrodites derived from *drsh-1/+* mothers, and in animals where maternal and zygotic Drosha was knocked down by RNAi (data not shown). Therefore, Drosha activity does not appear to be required for meiotic H3K9me2 accumulation.

We analyzed H3K9me2 accumulation in males carrying mutations in several other RNAi pathway components. RDE-1 (an Argonaute/Piwi/PAZ protein), RDE-2 (a novel protein), and RDE-4 (a dsRNA binding protein)

are required for RNAi upon introduction of exogenous dsRNA [19, 27]. RDE-2 is also required for normal germline development under conditions of stress [19, 28]. We observed normal H3K9me2 staining in male *rde-1*, *rde-2*, and *rde-4* mutants raised at both normal and elevated temperatures (data not shown). Hence, proteins required for RNAi per se are not required for the normal pattern of H3K9me2 accumulation during meiosis. *C. elegans* contains another 26 predicted Piwi/PAZ proteins (see Wormbase, <http://www.wormbase.org/>; [19]), several of which are known to function in RNAi-related phenomena. We observed normal H3K9me2 staining in *alg-1* (Argonaute-like gene required for microRNA function) [29] mutant males and upon RNAi knockdown of several other Piwi/PAZ candidates assayed thus far (data not shown). Systematic analysis of single and multiple Piwi/PAZ mutants is required to fully evaluate whether this protein family participates in meiotic heterochromatin assembly. Interestingly, RRF-1, DCR-1, and several RDE proteins have recently been reported to function in transcriptional silencing of a multicopy transgene in the *C. elegans* soma [30], although the H3K9 methylation status of repetitive transgenes in somatic cells is not known.

### *ego-1* Interacts Genetically with *him-17*

*ego-1* loss-of-function mutants have meiotic defects; leptotene-zygotene is protracted, and univalents are present in some oocytes at diakinesis (Figure 3C) [9, 10]. Univalents can result from defective chromosomal pairing, synaptonemal complex formation, or recombination [31]. Recent findings in yeast and *C. elegans* suggest that recombination relies on proper chromatin structure [13, 32]. Mutations in *C. elegans him-17* disrupt formation of double-strand breaks and lead to mei-






Figure 3. *ego-1* Interacts Genetically with *him-17*

Each panel shows one hermaphrodite gonad arm at 24 hr post-L4 stage. Asterisk indicates distal end of each arm. TZ, “transition zone” containing germ cells in early meiotic prophase (leptotene-zygotene stages). *ego-1* and *him-17* null alleles are shown (see [Supplemental Experimental Procedures](#)). Animals were raised at 20°C.

(A) The wild-type germline includes distal mitotic cells, meiotic cells, and oocytes in diakinesis stage of meiosis. Sperm are present at the proximal end (although most are out of the plane of focus).

(B) Organization of the *him-17* germline is normal.

(C) Overall organization of the *ego-1* germline is normal, although some nuclei are enlarged, the leptotene-zygotene region is expanded, and oocytes are small.

(D) The *ego-1;him-17* germline is small and lacks oocytes. Sperm nuclei are irregular in size (inset).

otic chromosome segregation defects [13]. They also severely reduce H3K9me2 accumulation on the male X chromosome and delay the transient H3K9me2 accumulation on hermaphrodite autosomes [13]. HIM-17 is a novel protein with several copies of the THAP domain implicated in DNA binding [13]. Since mutations in *ego-1* and *him-17* disrupt H3K9me2 enrichment of the male X chromosome and cause defects in early meiosis, we looked for a genetic interaction between them.

Wild-type *C. elegans* hermaphrodites produce sperm during the L4 larval stage and switch to oogenesis at the time of the adult molt. At 24 hr post-L4 stage, the wild-type hermaphrodite gonad contained 589 ( $\pm 21$ ;  $n = 2$ ) mitotic/early meiotic germ cells (including oocytes) and a reserve of >100 sperm (Figure 3A). In *him-17(0)* mutants raised at 20°C, germline function and organization are grossly normal, except for the appearance of univalent chromosomes in oocytes (Figure 3B) [13]. At 24 hr post-L4 stage, *him-17* mutants had 418 ( $\pm 2$ ;  $n = 2$ ) and *ego-1* mutants had 265 ( $\pm 40$ ;  $n = 8$ ) mitotic/early meiotic germ cells; >100 sperm were present in each case.

Adult *ego-1;him-17* hermaphrodites raised at 20°C have more severe germline defects compared with each single mutant (Figure 3D). At 24 hr post-L4 stage, the gonad arm contained only 50 ( $\pm 26$ ) sperm and 154 ( $\pm 34$ ) mitotic/early meiotic germ cells ( $n = 6$ ); oocytes were absent. Sperm nuclei were irregular in size and larger than normal, perhaps reflecting a combination of aneuploidy caused by chromosome segregation defects and improper chromosome condensation caused by a chromatin defect (Figure 3D, inset). By 48 hr post-L4 stage, only ~2% of *ego-1;him-17* germ lines contained any oocytes ( $n = 54$ ). The absence of oocytes may reflect either a sex determination defect (failure to make the sperm-to-oocyte switch) or the small germline (which does not produce sperm on schedule). *ego-1;him-17(0)* males also had small germlines and irregular sperm nuclei (data not shown).

We examined H3K9me2 accumulation in *ego-1;him-17* germlines to determine whether H3K9me2 levels were further reduced beyond either single mutant, but the abnormal chromosome morphology made it difficult to determine how the staining pattern was affected.




Figure 4. Two Models for the Role of EGO-1 in Silencing Unpaired DNA during Meiosis

(A and B) Unpaired DNA (A) associates with and (B) is transcribed by a DNA-dependent RNA polymerase that is not active on paired DNA. HIM-17 may participate in this process.

(C) EGO-1 binds to the RNA and recruits chromatin modifiers, e.g., histone methyltransferase (HMTase), to the region. RHA-1 may function in an RdRC-like complex with EGO-1 at this step. HMTase may directly bind to EGO-1 for its recruitment. Alternatively, EGO-1 generates dsRNA, which then triggers assembly of a RITS-like complex. RITS-like activity then recruits HMTase.

(D) The end result of the EGO-1-RNA association is accumulation of H3K9me2 modifications preferentially on the unpaired DNA.

The overall pattern appeared to be aberrant. A low level of H3K9me2 staining was seen, particularly in the middle and proximal end of the germline where meiotic nuclei are typically present, and some highly condensed nuclei were more brightly stained (data not shown).

#### EGO-1 Protein Is Enriched in the Nuclear Fraction of Cell Lysates

We used affinity-purified anti-EGO-1 polyclonal antibodies to assay for EGO-1 in nuclear extracts (see [Supplemental Experimental Procedures](#)). These antibodies do not work for immunofluorescence studies with a variety of fixation conditions, but they performed well in protein blots [11]. We prepared nuclear-enriched and total cellular protein extracts and performed Western blots with anti-EGO-1, anti-lamin (a nuclear control), and anti-myosin (a cytoplasmic control) antibodies, and we quantified the results. Most of the total cellular EGO-1 protein was detected in the nuclear extract, as would be expected if EGO-1 directly interacts with chromatin (Figure S3).

#### Role of EGO-1 in Meiotic Silencing

We find that EGO-1 activity is required for H3K9me2 enrichment on unsynapsed chromosomal regions during meiosis in *C. elegans*. This process appears to be dependent on an RdRP but independent of Dicer, and thus to be distinct from the mechanism of heterochromatin assembly described in *S. pombe* in which both RdRP and Dicer activities are required [3, 8]. Recognition/ repression of unpaired DNA in meiosis occurs in diverse species [3, 4] and may protect against spurious recombination and/or significant changes in the ordered sequences along homologs. Meiotic silencing

may also protect the germline from expression/movement of foreign DNA that has inserted into the genome and transposable genetic elements that have been mobilized. As such, meiotic silencing may complement RNA silencing mechanisms that repress the movement of transposable genetic elements via a posttranscriptional process [33].

It is unclear how unpaired DNA is recognized and repressed, and evidence suggests that different mechanisms may act in different species [3, 4]. Mammals use a chromatin-based mechanism that includes H3K9me2 and other histone modifications [4]. In contrast, *N. crassa* uses a mechanism that is thought to be posttranscriptional [3, 5]. Intriguingly, the *N. crassa* mechanism requires an RdRP, SAD-1, whose function is essential for meiosis. Perhaps EGO-1 participates in an ancient chromatin-based genome defense mechanism that has been adapted for essential germ cell processes. Indeed, gene expression profiling suggests that EGO-1 activity ultimately represses expression of genes whose activity may be detrimental to germline function and upregulates expression of genes whose activity is critical for germline function (V.E.V., V. Reinke, and E.M.M., unpublished data). It will be interesting to determine whether EGO-1 activity regulates the pattern of chromatin assembly at the repressed genes.

Although we detected only the gross level changes in H3K9me2 accumulation in *ego-1* germ cells, we expect that changes also occur at autosomal sites too scattered for detection by our methods. *ego-1* and *him-17* single mutants have subtly different defects in H3K9me2 accumulation (this study; [13]); therefore, EGO-1 and HIM-17 may regulate heterochromatin assembly at overlapping, but nonidentical, sets of chro-

mosomal sites. The *ego-1;him-17* phenotype may arise from the presence of a larger number of misregulated sites in the double mutant and/or additive defects from their nonoverlapping roles in other processes.

We propose two alternative models for how EGO-1 might target H3K9me2 to unpaired DNA (Figure 4). In both models, an RNA polymerase transcribes unpaired DNA, perhaps through unregulated initiation. EGO-1, in a complex with an RNA helicase (RHA-1) analogous to the RdRC complex in *S. pombe* [34], recognizes and binds the RNA as it is synthesized. RHA-1 is also required for H3K9me2 accumulation in *C. elegans* germ cells, but only at elevated temperatures, perhaps reflecting a purely structural role in this process [35]. In one model, EGO-1 converts ssRNA to dsRNA, which then triggers assembly of a RITS-type complex that promotes formation of H3K9me2 modifications, as proposed in other systems (see [8]). This model is supported by our finding that H3K9me2 accumulation is disrupted in *ego-1(RdRP/null)* putative catalytic mutants. The dsRNA may not require cleavage by a Dicer-type activity, perhaps reflecting a nonprocessive RdRP activity similar to that described for *N. crassa* QDE-1 in vitro [36]. In an alternative model, EGO-1 acts as a scaffold protein, bringing the chromatin-modifying machinery into proximity with unpaired DNA through its interaction with the transcript. There is ample precedent for the involvement of noncoding transcripts in facultative heterochromatin assembly, from X-inactivation to epigenetic imprinting [3, 4, 8, 37, 38], as well as recent evidence that transcription is required for targeting H3K9me2 to heterochromatic loci [39].

#### Supplemental Data

Supplemental Data include three figures and Supplemental Experimental Procedures and can be found with this article online at <http://www.current-biology.com/cgi/content/full/15/21/1972/DC1/>.

#### Acknowledgements

We thank C. David Allis, Jun Kelly Liu, and Henry Epstein for antibodies, Tim Schedl and Rodolfo Aramayo for discussion, Katherine Walstrom, Rob Martienssen, John Belote, and anonymous reviewers for comments on the manuscript, Claudia Zraly and Larry Smart for advice on methods, and Meera Sundaram, Shohei Mitani, and Geraldine Seydoux for strains. Some strains used in this study were provided by the *Caenorhabditis* Genetics Center, which is funded by the National Institutes of Health National Center for Research Resources. Funds for this study were provided by the National Science Foundation and Syracuse University (to E.M.M.) and the National Institutes of Health and Emory University Research Council (to W.G.K.).

Received: May 11, 2005

Revised: September 6, 2005

Accepted: September 23, 2005

Published: November 7, 2005

#### References

1. Bean, C.J., Schaner, C.E., and Kelly, W.G. (2004). Meiotic pairing and imprinted X chromatin assembly in *Caenorhabditis elegans*. *Nat. Genet.* 36, 100–105.
2. Kelly, W.G., Schaner, C.E., Dernberg, A.F., Lee, M.-H., Kim, S.K., Villeneuve, A.M., and Reinke, V. (2002). X-chromosome silencing in the germline of *C. elegans*. *Development* 129, 479–492.
3. Matzke, M.A., and Birchler, J.A. (2005). RNAi-mediated pathways in the nucleus. *Nat. Genet.* 6, 24–35.
4. Lee, J.T. (2005). Sex chromosome inactivation: the importance of pairing. *Curr. Biol.* 15, R249–R252.
5. Hynes, M.J., and Todd, R.B. (2003). Detection of unpaired DNA at meiosis results in RNA-mediated silencing. *Bioessays* 25, 99–103.
6. Ahlquist, P. (2002). RNA-dependent RNA polymerases, viruses, and RNA silencing. *Science* 296, 1268–1273.
7. Tijsterman, M., Ketting, R.F., and Plasterk, R.H.A. (2002). The genetics of RNA silencing. *Annu. Rev. Genet.* 36, 489–519.
8. Grewal, S.I.S., and Rice, J.C. (2004). Regulation of heterochromatin by histone methylation and small RNAs. *Curr. Opin. Cell Biol.* 16, 230–238.
9. Smardon, A., Spoerke, J.M., Stacey, S.C., Klein, M.E., Mackin, N., and Maine, E.M. (2000). EGO-1 is related to RNA-directed RNA polymerase and functions in germline development and RNA interference in *C. elegans*. *Curr. Biol.* 10, 169–178.
10. Qiao, L., Lissemore, J.L., Shu, P., Smardon, A., Gelber, M., and Maine, E.M. (1995). Enhancers of *glp-1*, a gene required for cell-signaling in *C. elegans*, define a set of genes required for germline development. *Genetics* 141, 551–569.
11. Vought, V.E., Ohmachi, M., Lee, M.-H., and Maine, E.M. (2005). EGO-1, a putative RNA-directed RNA polymerase, promotes germline proliferation in parallel with GLP-1/Notch signaling and regulates the spatial organization of nuclear pore complexes and germline P granules in *C. elegans*. *Genetics* 170, 1121–1132.
12. Simmer, F., Tijsterman, M., Parrish, S., Koushika, S.P., Nonet, M.L., Fire, A., Ahringer, J., and Plasterk, R.H.A. (2002). Loss of the putative RNA-directed RNA polymerase RRF-3 makes *C. elegans* hypersensitive to RNAi. *Curr. Biol.* 12, 1317–1319.
13. Reddy, K.C., and Villeneuve, A.M. (2004). *C. elegans* HIM-17 links chromatin modification and competence for initiation of meiotic recombination. *Cell* 118, 439–452.
14. Stinchcomb, D.T., Shaw, J.E., Carr, S.H., and Hirsh, D. (1985). Extrachromosomal DNA transformation of *Caenorhabditis elegans*. *Mol. Cell. Biol.* 5, 3484–3496.
15. Kelly, W.G., Xu, S., Montgomery, M.M., and Fire, A. (1997). Distinct requirements for somatic and germline expression of a generally expressed *Caenorhabditis elegans* gene. *Development* 146, 227–238.
16. Jedrusik, M.A., and Schulze, E. (2001). A single histone H1 isoform (H1.1) is essential for chromatin silencing and germline development in *Caenorhabditis elegans*. *Development* 128, 1069–1080.
17. Couteau, F., Guerry, F., Muller, F., and Palladino, F. (2002). A heterochromatin protein 1 homologue in *Caenorhabditis elegans* acts in germline and vulval development. *EMBO Rep.* 3, 235–241.
18. Kim, J.K., Gabel, H.W., Kamath, R.S., Tewari, M., Pasquinelli, A., Rual, J.F., Kennedy, S., Dybbs, M., Bertin, N., Kaplan, J.M., et al. (2005). Functional genomic analysis of RNA interference in *C. elegans*. *Science* 308, 1164–1167.
19. Tabara, H., Sarkissian, M., Kelly, W.G., Fleenor, J., Grishok, A., Timmons, L., Fire, A., and Mello, C.C. (1999). The *rde-1* gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* 99, 123–132.
20. Holdeman, R., Nehrt, S., and Strome, S. (1998). MES-2, a maternal protein essential for viability of the germline in *Caenorhabditis elegans*, is homologous to a *Drosophila* Polycomb group protein. *Development* 125, 2457–2467.
21. Bender, L.B., Cao, R., Zhang, Y., and Strome, S. (2004). THE MES-2/MES-3/MES-6 complex and regulation of histone H3 methylation in *C. elegans*. *Curr. Biol.* 14, 1639–1643.
22. Ketting, R.F., Haverkamp, T.H.A., van Leunen, H.G.A.M., and Plasterk, R.H.A. (1999). *mut-7* of *C. elegans*, required for transposon silencing and RNA interference, is a homolog of Werner Syndrome helicase and RNaseD. *Cell* 99, 133–141.
23. Robert, V.J.P., Sijen, T., van Wolfwinkel, J., and Plasterk, R.H.A. (2005). Chromatin and RNAi factors protect the *C. elegans* germline against repetitive sequences. *Genes Dev.* 19, 782–787.
24. Petrie, V.J., Wuitschick, J.D., Givens, C.D., Kosinski, A.M., and Partridge, J.F. (2005). RNA interference (RNAi)-dependent and RNAi-independent association of the Chp1 chromo-domain

- protein with distinct heterochromatic loci in fission yeast. *Mol. Cell. Biol.* **25**, 2331–2346.
25. Kanellopoulou, C., Muljo, S.A., Kung, A.L., Ganesan, S., Drapkin, R., Jenuwein, T., Livingston, D.M., and Rajewsky, K. (2005). Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev.* **19**, 489–501.
  26. Denli, A.M., Tops, B.B., Plasterk, R.H., Ketting, R.F., and Hannon, G.J. (2004). Processing of primary microRNAs by the Microprocessor complex. *Nature* **432**, 231–235.
  27. Tabara, H., Yigit, E., Siomi, H., and Mello, C.C. (2002). The dsRNA binding protein RDE-4 interacts with RDE-1, DCR-1, and a DExH-box helicase to direct RNAi in *C. elegans*. *Cell* **109**, 861–871.
  28. Tops, B.B., Tabara, H., Sijen, T., Simmer, F., Mello, C.C., Plasterk, R.H., and Ketting, R.F. (2005). RDE-2 interacts with MUT-7 to mediate RNA interference in *Caenorhabditis elegans*. *Nucleic Acids Res.* **33**, 347–355.
  29. Grishok, A., Pasquinelli, A.E., Conte, D., Li, N., Parrish, S., Ha, I., Baillie, D.L., Fire, A., Ruvkun, G., and Mello, C.C. (2001). Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing. *Cell* **106**, 23–34.
  30. Grishok, A., Sinskey, J.L., and Sharp, P.A. (2005). Transcriptional silencing of a transgene by RNAi in the soma of *C. elegans*. *Genes Dev.* **19**, 683–696.
  31. Page, S.L., and Hawley, R.S. (2003). Chromosome choreography: the meiotic ballet. *Science* **301**, 785–789.
  32. Maleki, S., and Keeney, S. (2004). Modifying histones and initiating meiotic recombination; new answers to an old question. *Cell* **118**, 404–406.
  33. Vastenhouw, N.L., and Plasterk, R.H. (2004). RNAi protects the *Caenorhabditis elegans* germline against transposition. *Trends Genet.* **20**, 314–319.
  34. Motamedi, M.R., Verdel, A., Colmenares, S.U., Gerber, S.A., Gygi, S.P., and Moazed, D. (2004). Two RNAi complexes, RITS and RDRC, physically interact and localize to noncoding centromeric RNAs. *Cell* **119**, 789–802.
  35. Walstrom, K.M., Schmidt, D., Bean, C.J., and Kelly, W.G. (2005). RNA helicase A is important for germline transcriptional control, proliferation, and meiosis in *C. elegans*. *Mech. Dev.* **122**, 707–720.
  36. Makeyev, E.V., and Bamford, D.H. (2002). Cellular RNA-dependent RNA polymerase involved in posttranscriptional gene silencing has two distinct activity modes. *Mol. Cell* **10**, 1417–1427.
  37. Craig, J. (2004). Heterochromatin—many flavours, common themes. *Bioessays* **27**, 17–28.
  38. Anderson, A.A., and Panning, B. (2003). Epigenetic gene regulation by noncoding RNAs. *Curr. Opin. Cell Biol.* **15**, 281–289.
  39. Schramke, V., Sheedy, D.M., Denli, A.M., Bonila, C., Ekwall, K., Hannon, G.J., and Allshire, R.C. (2005). RNA-interference-directed chromatin modification coupled to RNA polymerase II transcription. *Nature* **435**, 1275–1279.